



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

On the Early Development of Chimæra. BASHFORD DEAN.

Emphasis was laid on the similarity of the embryonic characters of Chimæroid and Elasmobranch.

Amphiuma and the Cæcilians. J. S. KINGSLEY.

The various statements which had been advanced to show the relationships of *Amphiuma* and *Cæcilians* were considered, and it was pointed out that these statements were almost entirely based upon misinterpretation or misconception. The differences between the two were then emphasized, and it was shown that the structural features were opposed to the view of Cope that the *Cæcilians* had descended from an *Amphiuma*-like form, and to that of the *Sarasins* that *Amphiuma* was a neoteric *Cæcilian*. In the possession of an ethmoid, in structure of vertebræ, in the relations of palatine and trigeminal nerves, in structure of nephridia and genitalia and in circulatory apparatus, the *Cæcilians* differ from *Amphiuma* and from all *Urodeles*, and the group must be regarded as entirely distinct from *Urodeles*, and as having descended directly from some *Stegocephalan* ancestor.

Vertebral Intercalation in Necturus. (Read by title.) H. C. BUMPUS.

Brachial and Lumbo-sacral Plexi in Necturus. F. C. WAITE.

In *Necturus maculosus* the normal position of the pelvic girdle is with attachment to the 19th vertebra, but in about one-fourth the cases it is attached to the 20th vertebra. Unfrequent cases are found in which the attachment is asymmetrical, the sacral rib on one side being one segment anterior to that on the other side.

Study of the plexi in a series of specimens shows: (a) that the position of the brachial plexus does not vary with displacement of pelvic girdle, and so it is im-

probable that intercalation of vertebræ occurs anterior to the posterior spinal nerve (V) involved in this plexus; (b) with normal position of pelvic girdle there are two prevalent types of topography of the lumbo-sacral plexus which depend upon the manner of branching of the spinal nerves to form the crural nerve, and further that there is considerable variation in the strength of the nerves involved, causing a shifting within narrow limits of the 'source center' of the plexus. (c) When the girdle is attached to 20th vertebra the plexus shows a displacement posteriorly, but not in a corresponding degree through an entire segment. It thus occupies a position intermediate between the normal position and what would be its position were it displaced through an entire segment. (d) Where the girdle is attached asymmetrically the plexus does not show corresponding asymmetry, but is essentially symmetrical in one of the two segments involved.

The intermediate position of the plexus, the occurrence of symmetrical variation in position of girdle; of asymmetrically placed girdles and of supernumerary sacral ribs, appears to be explicable not upon ground of intercalation of vertebra nor of slipping of girdle during ontogeny, but upon the hypothesis that there are several segments in this region, in any one of which a girdle may be developed.

Discovery of a Huge Octopus on the Coast of Florida. A. E. VERRILL.

The following officers were elected: President, C. S. Minot, Harvard; Vice-President, S. I. Smith, Yale; Secretary-Treasurer, G. H. Parker, Harvard; Members of the Executive Committee from the Society at large, J. S. Kingsley, Tufts, and Bashford Dean, Columbia.

G. H. PARKER,

HARVARD UNIVERSITY.

Secretary.